

NAME: _____

DATE: _____

MD: _____

TRANSDIAGNOSTIC GLOBAL IMPRESSION - PSYCHOPATHOLOGY

PART I.

The TGI-P is rated on a 7-point scale using a range of responses from 1 (normal) through to 7 (extreme severity).

After or during the visit, please rate the severity of symptoms considering your total clinical experience with patients:

	Normal	Minimal	Mild	Moderate	Marked	Severe	Extreme
1. Positive symptoms <i>Delusions, hallucinations, disorganised thinking, disorganised speech, abnormal motor behaviour</i>	1	2	3	4	5	6	7
2. Hostility symptoms <i>Anger, tension, uncooperativeness, impulsivity, aggression, irritability</i>	1	2	3	4	5	6	7
3. Manic symptoms <i>Expansive mood, grandiosity, racing thoughts, increased energy, excessive involvement in pleasurable activities</i>	1	2	3	4	5	6	7
4. Addiction symptoms <i>Impaired substance use control, craving, physical dependence</i>	1	2	3	4	5	6	7
5. Sleep symptoms <i>Hypersomnia or insomnia</i>	1	2	3	4	5	6	7
6. Negative symptoms <i>Blunted affect, alogia, asociality, avolition, anhedonia*</i>	1	2	3	4	5	6	7
7. Self-harm symptoms <i>Non-suicidal self-injury, suicidal ideation, intent, or attempt</i>	1	2	3	4	5	6	7
8. Depressive symptoms <i>Low mood, anhedonia, persistent feeling of sadness, hopelessness, helplessness</i>	1	2	3	4	5	6	7
9. Cognitive symptoms <i>Problems with concentration, attention, memory</i>	1	2	3	4	5	6	7
10. Anxiety symptoms <i>Feeling nervous, restless, tense, or fear of social interactions</i>	1	2	3	4	5	6	7

PART II.

To develop the TGI-P footprint, please insert the scores to the circles. First choose a symptom line and then put a dot where the line crosses the circle with the adequate severity. After rating every symptom, connect the dots to develop a symptom footprint. Use the TGI-P during every visit to compare changes and have a better understanding of the treatment progress.

